

---

# Twistor inspired developments in perturbative QCD

Nigel Glover

IPPP, University of Durham



SUSY 2005

The Millennium Window to Particle Physics

Durham, 20 July 2005

# How to calculate scattering amplitudes

---

- **Off-shell** methods  
Traditional Feynman diagram approach for off-shell  
Greens functions
- **On-shell** methods  
Based on S-matrix ideas of 1960's but recently inspired  
by **Witten's** proposal to relate **perturbative gauge  
theory amplitudes** to **topological string theory in  
twistor space**

Witten, hep-th/0312171

- ⇒ MHV rules
- ⇒ BCFW recursion relations
- ⇒ Generalised unitarity

- **Common methods**
  - Colour ordered amplitudes
  - Spinor helicity approach

# Common methods : Colour Ordered Amplitudes


---

$$\mathcal{A}_n(1, \dots, n) = \sum_{perms} Tr(T^{a_1} \dots T^{a_n}) A_n(1, \dots, n)$$

Colour-stripped amplitudes  $A_n$ : cyclically ordered

Order of external gluons fixed

The subamplitudes  $A_n$  have nice properties in the infrared limits.



Can reconstruct the full amplitude  $\mathcal{A}_n$  from  $A_n$ .  
In the large  $N$  limit,

$$|\mathcal{A}_n(1, \dots, n)|^2 \sim N^{n-2} \sum_{perms} |A_n(1, \dots, n)|^2$$

# Common methods : Spinor Helicity Formalism

---

- In Weyl (chiral) representation, each helicity state is represented by a bi-spinor ( $a = 1, 2$ )

$$\begin{aligned}u_+(p) &= \lambda_{pa}, & u_-(p) &= \tilde{\lambda}_p^{\dot{a}}, \\ \overline{u_+(p)} &= \tilde{\lambda}_{p\dot{a}}, & \overline{u_-(p)} &= \lambda_p^a\end{aligned}$$

so that

$$\begin{aligned}\langle ij \rangle &= \overline{u_-(p_i)} u_+(p_j) = \lambda_i^a \lambda_{ja} = \epsilon_{ab} \lambda_i^a \lambda_j^b \\ [ij] &= \overline{u_+(p_i)} u_-(p_j) = \tilde{\lambda}_{i\dot{a}} \tilde{\lambda}_j^{\dot{a}} = -\epsilon_{\dot{a}\dot{b}} \tilde{\lambda}_i^{\dot{a}} \tilde{\lambda}_j^{\dot{b}}\end{aligned}$$

- We can write massless vector

$$p_{a\dot{a}} \equiv p_\mu \sigma_{a\dot{a}}^\mu = \lambda_{pa} \tilde{\lambda}_{p\dot{a}}$$

# Common methods : Spinor Helicity Formalism

---

- Polarisation vectors for particle  $i$ :

$$\epsilon_{ia\dot{a}}^- = \frac{\lambda_{ia} \tilde{\eta}_{\dot{a}}}{[\tilde{\lambda}_i \tilde{\eta}]}, \quad \epsilon_{ia\dot{a}}^+ = \frac{\eta_a \tilde{\lambda}_{i\dot{a}}}{\langle \eta \lambda_i \rangle}$$

- For **real** momenta in Minkowski space,

$$\tilde{\lambda} = \lambda^*$$

- For space-time signature  $(+, +, -, -)$ ,  $\tilde{\lambda}, \lambda$  are real and independent
- Amplitudes are functions of the  $\lambda_i$  and  $\tilde{\lambda}_i$

# Off-shell methods: Feynman diagrams

---

- ✓ **Direct link to Lagrangian**
- ✓ **Easy to adapt to any model**
- ✓ **Easy to include massive particles with/without spin**
- ✓ **Easy to automate**  
⇒ **tree-level packages** Madgraph/Grace/CompHep/...
- ✓ **Off-shell Berends-Giele recursion relations**  
⇒ **tree-level packages** AlpGen/HELAC/PHEGAS/...
- ✗ **Many Feynman diagrams**
- ✗ **Large cancellations between diagrams**
- ✗ **Loop amplitudes manpower intensive**

# Example

---


## Multi-jet production at the LHC using HELAC/PHEGAS

Draggiotis, Kleiss, Papadopoloulos

# of jets	2	3	4	5	6	7	8
# of dist.processes	10	14	28	36	64	78	130
total # of processes	126	206	621	861	1862	2326	4342
$\sigma(nb)$	-	91.41	6.54	0.458	0.030	0.0022	0.00021
% Gluonic	-	45.7	39.2	35.7	35.1	33.8	26.6

- The number of Feynman diagrams for an  $n$  gluon process increases very quickly with  $n$
- ⇒ for the 10 gluon amplitude there are 10,525,900 diagrams
- ⇒ Feynman diagrams very inefficient for many legs
- Control the quantum numbers of the scattering particles

# Gluonic helicity amplitudes



Each row describes scattering with  $n_+$  positive helicities and  $n_-$  negative helicities.

Each circle represents an allowed helicity configuration - from all positive on the right to all negative on the left



# Gluonic helicity amplitudes

---

For example, the result of computing the 25 diagrams for the five-gluon process yields

$$A_5(1^\pm, 2^+, 3^+, 4^+, 5^+) = 0$$

$$A_5(1^-, 2^-, 3^+, 4^+, 5^+) = \frac{\langle 12 \rangle^4}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 45 \rangle \langle 51 \rangle}$$

In fact, for  $n$  point amplitudes,


$$A_n(1^\pm, 2^+, 3^+, \dots, n^+) = 0$$

$$A_n(1^-, 2^-, 3^+, \dots, n^+) = \frac{\langle 12 \rangle^4}{\langle 12 \rangle \langle 23 \rangle \cdots \langle n1 \rangle}$$

Maximally helicity violating (MHV) amplitudes

Parke, Taylor; Berends, Giele


# Gluonic helicity amplitudes



$$A_n(1^\pm, 2^+, 3^+, \dots, n^+) = 0$$

effective tree-level supersymmetry

# Gluonic helicity amplitudes



$$A_n(1^-, 2^-, 3^+, \dots, n^+) = \frac{\langle 12 \rangle^4}{\langle 12 \rangle \langle 23 \rangle \cdots \langle n1 \rangle}$$

# Twistor Space

---

Penrose, 1967

Amplitudes in twistor space obtained by Fourier transform with respect to positive helicity spinors,

$$\tilde{\lambda}_{\dot{a}} = i \frac{\partial}{\partial \mu^{\dot{a}}}, \quad \mu^{\dot{a}} = i \frac{\partial}{\partial \tilde{\lambda}_{\dot{a}}}$$

Momentum conservation yields

$$\delta \left( \sum k_i \right) = \int d^4x \exp \left( i x^{a\dot{a}} \sum_j \lambda_{j\dot{a}} \tilde{\lambda}_{j\dot{a}} \right)$$

so that the amplitude in twistor space is

$$\tilde{A}(\lambda_i, \mu_i) = \int d^4x \int \prod_i \frac{d^2 \tilde{\lambda}_i}{(2\pi)^2} \exp \left( i \sum_j (\mu_j^{\dot{a}} + x^{a\dot{a}} \lambda_{j\dot{a}}) \tilde{\lambda}_{j\dot{a}} \right) A(\lambda_i, \tilde{\lambda}_i)$$

# Twistor Space


---

Witten, hep-th/0312171

Witten observed that in twistor space external points lie on certain algebraic curves



⇒ degree of curve is related to the number of negative helicities and loops

$$d = n_- - 1 + l$$





# Twistor Space

MHV



NMHV




# MHV rules

Start from **on-shell** MHV amplitude and define **off-shell** vertices


Cachazo, Svrcek and Witten

$$V(1^-, 2^-, 3^+, \dots, n^+, P^+) = \frac{\langle 12 \rangle^4}{\langle 12 \rangle \dots \langle n-1n \rangle \langle nP \rangle \langle P1 \rangle}$$



and

$$V(1^-, 2^+, 3^+, \dots, n^+, P^-) = \frac{\langle 1P \rangle^4}{\langle 12 \rangle \dots \langle n-1n \rangle \langle nP \rangle \langle P1 \rangle}$$



Crucial step is **off-shell** continuation  $P^2 \neq 0$ :


$$\langle iP \rangle = \frac{\langle i^- | \not{P} | \eta^- \rangle}{[P\eta]} = \sum_j \frac{\langle i^- | j | \eta^- \rangle}{[P\eta]}$$

where  $P = \sum_j j$  and  $\eta$  is lightlike auxiliary vector

# MHV rules

---

Must connect up a positive helicity off-shell line to a negative helicity off-shell line with a scalar propagator



Connecting two MHV's  $\Rightarrow$  amplitude with 3 negative helicities

Connecting three MHV's  $\Rightarrow$  amplitude with 4 negative helicities

etc.



# Example: six gluon scattering

---


As an example, let's use the MHV rules to calculate one of the first non-MHV amplitudes

$$A_6(1^-, 2^-, 3^-, 4^+, 5^+, 6^+)$$

**Step 1** Draw all the allowed MHV diagrams





# Example: six gluon scattering

There are six MHV graphs



# Example: six gluon scattering

Some graphs are not allowed e.g.



# Example: six gluon scattering

---



As an example, let's use the MHV rules to calculate one of the first non-MHV amplitudes

$$A_6(1^-, 2^-, 3^-, 4^+, 5^+, 6^+)$$

**Step 1** Draw all the allowed MHV diagrams

**Step 2** Apply MHV rules to each diagram


# Example: six gluon scattering: diagram 1



$$\frac{\langle 12 \rangle^4}{\langle 56 \rangle \langle 61 \rangle \langle 12 \rangle \langle 2|P|\eta \rangle \langle 5|P|\eta \rangle} \times \frac{1}{s_{34}} \times \frac{\langle 3|P|\eta \rangle^4}{\langle 34 \rangle \langle 4|P|\eta \rangle \langle 3|P|\eta \rangle}$$

with  $P = 3 + 4 = -(1 + 2 + 5 + 6)$

# Example: six gluon scattering: diagram 2



$$\frac{\langle 12 \rangle^4}{\langle 61 \rangle \langle 12 \rangle \langle 2|P|\eta \rangle \langle 6|P|\eta \rangle} \times \frac{1}{s_{345}} \times \frac{\langle 3|P|\eta \rangle^4}{\langle 34 \rangle \langle 45 \rangle \langle 5|P|\eta \rangle \langle 3|P|\eta \rangle}$$

with  $P = 3 + 4 + 5 = -(1 + 2 + 6)$

# Example: six gluon scattering

---

As an example, let's use the MHV rules to calculate one of the first non-MHV amplitudes

$$A_6(1^-, 2^-, 3^-, 4^+, 5^+, 6^+)$$


- Step 1** Draw all the allowed MHV diagrams
- Step 2** Apply MHV rules to each diagram
- Step 3** Add up diagrams and check  $\eta$  independence

# Next-to MHV amplitude for $n$ gluons

Simplest case:  $A_n(1^-, 2^-, 3^-, 4^+, \dots, n^+)$

$2(n - 3)$  graphs

Cachazo, Svrcek and Witten



$$\begin{aligned}
 A &= \sum_{i=3}^{n-1} \frac{\langle 1|(2, i)|\eta\rangle^3}{\langle (i+1)|(2, i)|\eta\rangle \langle i+1 i+2\rangle \dots \langle n1\rangle} \frac{1}{s_{2,i}^2} \frac{\langle 23\rangle^3}{\langle 2|(2, i)|\eta\rangle \langle 34\rangle \dots \langle i|(2, i)|\eta\rangle} \\
 &+ \sum_{i=4}^n \frac{\langle 12\rangle^3}{\langle 2|(3, i)|\eta\rangle \langle (i+1)|(3, i)|\eta\rangle \dots \langle n1\rangle} \frac{1}{s_{3,i}^2} \frac{\langle 3|(3, i)|\eta\rangle^3}{\langle 34\rangle \dots \langle i-1 i\rangle \langle i|(3, i)|\eta\rangle}.
 \end{aligned}$$


where  $(k, i) = k + \dots + i$  is the off-shell momentum

$\Rightarrow$  Lorentz invariant and gauge invariant expressions



# Generating all the tree amplitudes

Amplitudes with  $i^-$  and  $j^+$  helicities



- MHV rules always adds one negative helicity and any number of positive helicities  
⇒ maps out all allowed tree amplitudes

# Other processes

---

MHV rules have been generalised to many other processes

✓ with massless fermions - quarks, gluinos

Georgiou and Khoze; Wu and Zhu; Georgiou, EWNG and Khoze

✓ with massless scalars - squarks

Georgiou, EWNG and Khoze; Khoze

✓ with an external Higgs boson

Dixon, EWNG, Khoze; Badger, EWNG, Khoze

✓ with an external weak boson

Bern, Forde, Kosower and Mastrolia

Has provided **new** analytic results for  $n$ -particle amplitudes


Also useful for studying infrared properties of amplitudes

Birthwright, EWNG, Khoze and Marquard

# BCFW on-shell recursion relations

Britto, Cachazo, Feng; Roiban, Spradlin, Volovich

Lets consider an  $n$  particle amplitude  $A(0)$ .



hatted momenta are shifted to put on-shell

$$\hat{i} = i + z\eta, \quad \hat{j} = j - z\eta, \quad \hat{P} = P + z\eta$$

⇒ each vertex is an **on-shell** amplitude

# BCFW recursion relations

---

- It turns out that the shift  $\eta$  is not a momentum, but

$$\eta = \lambda_i \tilde{\lambda}_j \quad OR \quad \eta = \lambda_j \tilde{\lambda}_i$$

- The parameter  $z$  is fixed by  $\hat{P}^2 = 0$

$$z = \frac{P^2}{\langle j|P|i \rangle}$$


- Easy to prove that by complex analysis based on fact that only simple poles in  $z$  occur and that  $A(z)$  vanishes as  $z \rightarrow \infty$

**Britto, Cachazo, Feng and Witten**

- Requires on-shell three-point vertex contributions - both MHV and  $\overline{\text{MHV}}$  .

# BCFW - six gluon example

If we select 3 and 4 to be the special gluons, there are only three diagrams (for any helicities)



For this helicity assignment, the middle diagram is zero!

$$A_6(1^-, 2^-, 3^-, 4^+, 5^+, 6^+)$$

$$= \frac{1}{\langle 5|\cancel{3} + \cancel{4}|2\rangle} \left( \frac{\langle 1|\cancel{2} + \cancel{3}|4\rangle^3}{[23][34]\langle 56\rangle\langle 61\rangle s_{234}} + \frac{\langle 3|\cancel{4} + \cancel{5}|6\rangle^3}{[61][12]\langle 34\rangle\langle 45\rangle s_{345}} \right)$$

Extremely compact analytic results for up to 8 gluons

# Other processes

---

BCF recursion relations have been generalised to other processes

- ✓ with massless fermions - quarks, gluinos

Luo and Wen

- ✓ gravitons

Bedford, Brandhuber, Spence and Travaglini; Cachazo and Svrcek

There is nothing (in principle) to stop this approach being applied to particles with mass.

- ✓ massive coloured scalars

Badger, EWNG, Khoze and Svrcek

- ✓ massive vector bosons and heavy quarks

Badger, EWNG and Khoze

# One loop amplitudes

---

- So far, supersymmetry was not a major factor - tree level amplitudes same for  $\mathcal{N} = 4$   $\mathcal{N} = 1$  and QCD
- Not true at the loop level due to circulating states


$$\begin{aligned}A_n^{\mathcal{N}=4} &= A_n^{[1]} + 4A_n^{[1/2]} + 3A_n^{[0]} \\A_n^{\mathcal{N}=1, \text{chiral}} &= A_n^{[1/2]} + A_n^{[0]} \\A_n^{\text{glue}} &= A_n^{\mathcal{N}=4} - 4A_n^{\mathcal{N}=1, \text{chiral}} + A_n^{[0]}\end{aligned}$$

- All plus and nearly all-plus amplitudes do not vanish for non-supersymmetric QCD
- A lot of progress by a lot of people

# One loop amplitudes

---

- Key point is that loop amplitudes contain **both poles** and **cuts** - e.g.  $\log(x)$  has cut for negative  $x$
- **Cut** contributions are constructible by using **unitarity** - Cut lines are **on-shell** and 4-dimensional



Bern, Dixon, Dunbar, Kosower; Britto, Cachazo, Feng

- **Pole** contributions are amenable to adapted BCFW recursion relations

Bern, Dixon, Kosower



# SUSY QCD loops

---

- ✓  $\mathcal{N} = 4$  and  $\mathcal{N} = 1$  one-loop amplitudes are constructible from their 4-dimensional cuts  
⇒ employ unitarity techniques

Bern, Dixon, Dunbar, Kosower

- ✓ For  $\mathcal{N} = 4$  **all** amplitudes are a linear combination of known box integrals


$$A_n = \Sigma \quad \mathbf{a} \quad \mathbf{b} \quad \mathbf{c} \quad \mathbf{d} \quad \mathbf{e} \quad \mathbf{f}$$

The image shows six Feynman diagrams representing box integrals, arranged in two rows of three. Each diagram is a square with four external lines. The diagrams are labeled a through f. Diagram a has red lines on the top-left and bottom-right corners. Diagram b has red lines on the top-right and bottom-left corners. Diagram c has red lines on the top-left and bottom-right corners. Diagram d has red lines on the top-left and bottom-right corners. Diagram e has red lines on the top-left and bottom-right corners. Diagram f has red lines on the top-left and bottom-right corners.

# Twistor space interpretation

- Coefficients of boxes have very interesting structures.

Bern, Del Duca, Dixon, Kosower; Britto, Cachazo, Feng





# Twistor space interpretation

---

- Four mass box first appears in eight-point amplitude with four negative and four positive helicities

Bern, Dixon, Kosower

e.g.



# QCD loops

---

QCD amplitudes more complicated

- (a) Not 4-dimensional cut constructible. Rational function contribution not probed by 4-d cut
- (b) All plus and almost all plus amplitudes not zero - but rational functions. Not protected by SWI.

Nevertheless, all four-point and five-point amplitudes known:

Recent progress

- ✓ On-shell recurrence relations for all plus and almost all plus amplitudes

Bern, Dixon and Kosower, hep-ph/0501240 and 0505055

Recursion relations complicated by double pole terms and boundary terms

- ✓ Rational parts of infrared divergent amplitudes computed using **on-shell** recursion relation

Bern, Dixon and Kosower, hep-ph/0507005

# Summary

---

- **On-shell** techniques are a very exciting and rapidly developing field
- **MHV** rules for tree-level  
Very simple way of deriving  $n$ -point amplitudes for massless partons
- **BCFW** recursion relations for tree-level  
Very powerful method for deriving amplitudes for both massless and massive particles

Badger, EWNG, Khoze and Svrcek


- **Generalised unitarity** and one-loop amplitudes  
**SUSY** amplitudes cut constructible - coefficients of loop integrals can be *read off* from graphs  
**QCD** amplitudes contain cut-non constructible parts. These simple pole terms can be attacked using the BCFW relations

Bern, Dixon, Kosower

Expect all one-loop six-point gluon amplitudes soon

# Berends-Giele : Off-shell recursion relations

Full amplitudes can be built up from simpler amplitudes with fewer particles



Purple gluons are off-shell, green gluons are on-shell.  
This is a recursion relation built from off-shell currents.

**Berends, Giele**

Particularly suited to numerical solution

ALPGEN, HELAC/PHEGAS